

A noncommutative version of Beilinson's Theorem

Hiroyuki Minamoto

February 2, 2008

Abstract

We prove that the category of representations of the N -Kronecker quiver and the category $\text{qgr}\left(k\langle X_1, \dots, X_N \rangle / (\sum_{i=1}^N X_i^2)\right)$ which is considered as the category of coherent sheaves on the noncommutative projective scheme associated to $k\langle X_1, \dots, X_N \rangle / (\sum_{i=1}^N X_i^2)$ are derived equivalent. This equivalence is easily proved by applying Orlov's Theorem, on the other hand, in our proof, the quadratic relation $\sum_{i=1}^N X_i^2$ naturally arises from Auslander-Reiten theory.

0 Introduction

In [Be] Beilinson proved that the derived category of $\mathbb{P}^n$ is generated by the line bundles $\mathcal{O}_{\mathbb{P}^n}, \dots, \mathcal{O}_{\mathbb{P}^n}(n)$. This is equivalent to saying that the functor

$$\mathbb{R} \text{Hom}\left(\bigoplus_{i=0}^n \mathcal{O}_{\mathbb{P}^n}(i), -\right) : D^b(\text{coh } \mathbb{P}^n) \longrightarrow D^b(\text{mod-}B)$$

is an equivalence of triangulated categories where $B = \text{End}(\bigoplus_{i=0}^n \mathcal{O}_{\mathbb{P}^n}(i))$. (This equivalence is also called Beilinson's Theorem.)

Although the projective space $\mathbb{P}_k^n$ is a basic object in algebraic geometry, the corresponding finite dimensional algebra B is not elementary:

B is the path algebra $k\vec{Q}_{n+1}$ of the Beilinson quiver $\vec{Q}_{n+1}$

$$\begin{array}{ccccccc} 0 & \xrightarrow{x_1^{(0)}} & 1 & \xrightarrow{x_1^{(1)}} & 2 & \xrightarrow{x_1^{(2)}} & \cdots \cdots \cdots \xrightarrow{x_1^{(n-2)}} & n-1 & \xrightarrow{x_1^{(n-1)}} & n \\ & \vdots & & \vdots & & \vdots & & \vdots & & \vdots \\ & x_{n+1}^{(0)} & & x_{n+1}^{(1)} & & x_{n+1}^{(2)} & & x_{n+1}^{(n-2)} & & x_{n+1}^{(n-1)} \end{array}$$

with the relations

$$x_i^{(l+1)} x_j^{(l)} = x_j^{(l+1)} x_i^{(l)} \quad \text{for } i, j \in \{1, \dots, n\} \text{ and } l = 0, \dots, n-2.$$

In this paper we consider the path algebra $k\overrightarrow{\Omega}_N$ of N -Kronecker quiver $\overrightarrow{\Omega}_N$

$$0 \begin{array}{c} \xrightarrow{x_1} \\ \vdots \\ \xrightarrow{x_N} \end{array} 1$$

which is one of the simplest non-trivial finite dimensional algebra. Our main theorem states that in noncommutative projective algebraic geometry, there is an imaginary geometric object $\text{proj } R$ which is derived equivalent to $\text{mod-}k\overrightarrow{\Omega}_N$.

The main result of this paper is the following.

Theorem 0.1 (Theorem 2.15). *Let k be an algebraically closed field, and $A = k\overrightarrow{\Omega}_N$, the path algebra of the N -Kronecker quiver $\overrightarrow{\Omega}_N$.*

Set $R = k\langle X_1, \dots, X_N \rangle / (\sum_{i=1}^N X_i^2)$. Then for $N \geq 2$, there exists a natural equivalence of derived categories

$$D^b(\text{mod-}A) \cong D^b(\text{qgr } R),$$

where $\text{mod-}A$ is the category of finite A -modules and $\text{qgr } R$ is the category which is considered as the category of coherent sheaves on the noncommutative projective scheme $\text{proj } R$ associated to the graded ring R .

This theorem is easily proved by applying Orlov's Theorem [O, Theorem 2.5]. Orlov's Theorem gives a sufficient condition for a graded ring R that $\text{proj } R$ is derived equivalent to $\text{mod-}k\overrightarrow{\Omega}_N$. But in contrast, our theorem states that for the path algebra $k\overrightarrow{\Omega}_N$ there is a canonical choice of such R .

To prove Theorem 0.1, we use a method of noncommutative projective geometry and a result on the derived Picard group by J. Miyachi and A. Yekutieli given in [MY]. They computed the derived Picard groups of finite dimensional path algebras of quivers by using Happel's derived version of Auslander-Reiten theory, from which the quadratic relation $\sum_{i=1}^N X_i^2$ naturally arises.

In [KR], Kontsevich and Rosenberg constructed the category Spaces_k of "noncommutative spaces," and showed that in the category Spaces_k , the functor represented by projective space $\mathbb{P}_k^n$ in (commutative) algebraic geometry is represented by an object $\text{N}\mathbb{P}_k^n$, the noncommutative projective space. They also proved that the category $\text{coh N}\mathbb{P}_k^n$ of coherent sheaves on $\text{N}\mathbb{P}_k^n$ is derived equivalent to the category $\text{mod-}k\overrightarrow{\Omega}_{n+1}$. Therefore it is natural to ask whether the graded ring $k\langle X_1, \dots, X_{n+1} \rangle / (\sum_{i=1}^{n+1} X_i^2)$ is the coordinate ring of $\text{N}\mathbb{P}_k^n$, i.e., whether $\text{coh N}\mathbb{P}_k^n$ is equivalent to $\text{qgr } k\langle X_1, \dots, X_{n+1} \rangle / (\sum_{i=1}^{n+1} X_i^2)$.

We organize the present paper as follows: in Section 1 we introduce some definitions and results: in Section 2 we prove Theorem 0.1 (Theorem 2.15).

Acknowledgment. I am grateful to I. Mori for his great help and comment on the first version of the paper. I am also grateful to T. Abe for his encouragement and A. Moriwaki for his hospitality and kindness. I thank A. Takahashi for pointing out the paper of Orlov [O].

1 Preliminaries

In this section we introduce some definitions and results. Some of them are about noncommutative projective schemes and the others are about derived Picard groups.

1.1 Noncommutative Projective Schemes

This subsection is a summary of the paper [Po] by A. Polishchuk. We only treat $\mathbb{N}$ -graded algebras although $\mathbb{Z}$ -algebras which is the more general notion are treated in [Po].

Let k be a field and let $R = k \oplus R_1 \oplus R_2 \oplus \cdots$ be a connected graded coherent ring. $\text{Gr } R$ (resp. $\text{gr } R$) denote the category of graded right R -modules (resp. finitely presented graded right R -modules). $\text{Tor } R$ (resp. $\text{tor } R$) denote the full subcategory of torsion modules (resp. modules finite dimensional over k). Note that $\text{Tor } R$ and $\text{tor } R$ are dense subcategories of $\text{Gr } R$ and $\text{gr } R$ respectively, hence the quotient categories $\text{QGr } R = \text{Gr } R / \text{Tor } R$ and $\text{qgr } R = \text{gr } R / \text{tor } R$ are abelian categories.

For graded a right R -module $M = \bigoplus_{n \in \mathbb{Z}} M_n$, we denote by $M(1)$ the 1-degree shift of M . i.e., $M(1)_n = M_{n+1}$. The degree shift operator $(1) : \text{gr } R \longrightarrow \text{gr } R$ induces the autoequivalence (1) on $\text{qgr } R$. We denote by $\overline{R}$ the image in $\text{qgr } R$ of the regular module R_R . The (coherent) *noncommutative projective scheme* $\text{proj } R$ associated to R is the triple $(\text{qgr } R, \overline{R}, (1))$. The autoequivalence (1) is called the *canonical polarization* on $\text{proj } R$.

In noncommutative projective geometry, one thinks of $\text{qgr } R$ as the category of coherent sheaves on a noncommutative projective scheme $\text{proj } R$ associated to a graded ring R ([AZ, SvB, Po]).

Let us consider a triple $(\mathcal{C}, \mathcal{O}, s)$ consisting of a k -linear abelian category $\mathcal{C}$, an object $\mathcal{O}$ and an autoequivalence s on $\mathcal{C}$. For $\mathcal{F} \in \mathcal{C}$, we define

$$\Gamma_*(\mathcal{F}) = \bigoplus_{n \geq 0} \text{Hom}_{\mathcal{C}}(\mathcal{O}, \mathcal{F}(n)),$$

where $\mathcal{F}(n) = s^n \mathcal{F}$, and we set

$$R = \Gamma_*(\mathcal{C}, \mathcal{O}, s) = \Gamma_*(\mathcal{O}).$$

Multiplication is defined as follows: If $x \in \text{Hom}_{\mathcal{C}}(\mathcal{O}, \mathcal{F}(l))$, $b \in \text{Hom}_{\mathcal{C}}(\mathcal{O}, \mathcal{O}(m))$ and $a \in \text{Hom}_{\mathcal{C}}(\mathcal{O}, \mathcal{O}(n))$ then

$$x \cdot a = s^n(x) \circ a \quad \text{and} \quad a \cdot b = s^m(a) \circ b.$$

With this law of composition, $\Gamma_*(\mathcal{F})$ becomes a graded right module over the graded algebra R over k .

Definition 1.1 ([AZ, Section 4.2], [Po, Section 2]). *Let $(\mathcal{C}, \mathcal{O}, s)$ be a triple as above. Then the pair $(\mathcal{O}, s)$ is called ample if the following conditions hold:*

- (1) For every object $\mathcal{F} \in \mathcal{C}$, there are positive integers $l_1, \dots, l_p$ and an epimorphism $\oplus_{i=1}^p \mathcal{O}(-l_i) \longrightarrow \mathcal{F}$.
- (2) For every epimorphism $f : \mathcal{F} \longrightarrow \mathcal{G}$, there exists an integer n_0 such that for every $n \geq n_0$ the induced map $\mathrm{Hom}_{\mathcal{C}}(\mathcal{O}, \mathcal{F}(n)) \longrightarrow \mathrm{Hom}_{\mathcal{C}}(\mathcal{O}, \mathcal{G}(n))$ is surjective.

Let $\pi : \mathrm{Gr} R \longrightarrow \mathrm{QGr} R$ be the quotient functor. Set $\bar{\Gamma}_* = \pi \circ \Gamma_*$.

Theorem 1.2 ([Po, Theorem 2.4]). *Let $(\mathcal{C}, \mathcal{O}, s)$ be a triple as above. If $(\mathcal{O}, s)$ is ample, then the graded ring $R = \Gamma_*(\mathcal{C}, \mathcal{O}, s)$ is coherent, $\Gamma_*(\mathcal{F})$ is finitely presented R -module for $\mathcal{F} \in \mathcal{C}$ and the functor $\bar{\Gamma}_* : \mathcal{C} \longrightarrow \mathrm{qgr} R$ induces an equivalence of triples between $(\mathcal{C}, \mathcal{O}, s)$ and $\mathrm{proj} R = (\mathrm{qgr} R, \bar{R}, (1))$, i.e.,*

$\bar{\Gamma}_* : \mathcal{C} \xrightarrow{\sim} \mathrm{qgr} R$ is an equivalence of categories,

$\bar{\Gamma}_*(\mathcal{O}) \cong \bar{R}$, and $\bar{\Gamma}_* \circ s = (1) \circ \bar{\Gamma}_*$.

By Beilinson's Theorem $\mathrm{coh} \mathbb{P}^1$ is derived equivalent to $\mathrm{mod}\text{-}k\vec{\Omega}_2$. Therefore by Theorem 0.1, $\mathrm{coh} \mathbb{P}^1$ and $\mathrm{qgr} \left(\frac{k\langle X_1, X_2 \rangle}{(X_1^2 + X_2^2)} \right)$ are derived equivalent. However, it turns out that these two categories are equivalent.

Example 1.3 ([SvB, Section 3]). *Let $\sigma : \mathbb{P}^1 \longrightarrow \mathbb{P}^1$, $[a : b] \mapsto [b : -a]$ be the automorphism of $\mathbb{P}^1$ and set $s = \sigma_*(- \otimes_{\mathcal{O}_{\mathbb{P}^1}} \mathcal{O}_{\mathbb{P}^1}(1))$. Then s is ample on the triple $(\mathrm{coh}(\mathbb{P}^1), \mathcal{O}_{\mathbb{P}^1}, s)$ and*

$$\Gamma_*(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}, s) \cong \frac{k\langle X_1, X_2 \rangle}{(X_1^2 + X_2^2)}$$

Traditionally the graded ring $\Gamma_*(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}, s)$ is denoted by $B(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(1), \sigma)$ and is called the twisted homogeneous coordinate ring associated to the pair $(\mathcal{O}_{\mathbb{P}^1}(1), \sigma)$.

Hence by Theorem 1.2, the triples $(\mathrm{coh} \mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}, s)$ and $\mathrm{proj} \left(\frac{k\langle X_1, X_2 \rangle}{(X_1^2 + X_2^2)} \right)$ are equivalent. In particular there is an equivalence

$$\bar{\Gamma}_* : \mathrm{coh} \mathbb{P}^1 \xrightarrow{\sim} \mathrm{qgr} \left(\frac{k\langle X_1, X_2 \rangle}{(X_1^2 + X_2^2)} \right).$$

Moreover, from the proof of Theorem 0.1 in Section 2, it is easy to see that there is a commutative diagram

$$\begin{array}{ccccc} D^b(\mathrm{coh} \mathbb{P}^1) & \xrightarrow{\sim} & D^b(\mathrm{mod}\text{-}k\vec{\Omega}_2) & \xleftarrow{\sim} & D^b \left(\mathrm{qgr} \left(\frac{k\langle X_1, X_2 \rangle}{(X_1^2 + X_2^2)} \right) \right) \\ \uparrow & & & & \uparrow \\ \mathrm{coh} \mathbb{P}^1 & \xrightarrow[\bar{\Gamma}_*]{\sim} & & & \mathrm{qgr} \left(\frac{k\langle X_1, X_2 \rangle}{(X_1^2 + X_2^2)} \right) \end{array}$$

where the vertical arrows are natural inclusions.

Remark 1.4. *The triples $(\text{coh } \mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}, - \otimes_{\mathbb{P}^1} \mathcal{O}_{\mathbb{P}^1}(1))$ and $\text{proj}(k[X_1, X_2])$ are equivalent by Serre's Theorem [Har, Proposition II.5.15]. Hence the categories $\text{qgr}(k[X_1, X_2])$ and $\text{qgr}\left(\frac{k\langle X_1, X_2 \rangle}{(X_1^2 + X_2^2)}\right)$ are equivalent.*

Moreover these two (noncommutative) projective schemes are equivalent at the level of affine covers. There is an equivalence

$$\text{gr}(k[X_1, X_2]) \xrightarrow{\sim} \text{gr}\left(\frac{k\langle X_1, X_2 \rangle}{(X_1^2 + X_2^2)}\right)$$

which sends $k[X_1, X_2]$ to $\frac{k\langle X_1, X_2 \rangle}{(X_1^2 + X_2^2)}$. As tor is the full subcategory of gr consisting of objects of finite length, from the above equivalence of affine covers we can also obtain the equivalence between $\text{qgr}(k[X_1, X_2])$ and $\text{qgr}\left(\frac{k\langle X_1, X_2 \rangle}{(X_1^2 + X_2^2)}\right)$. (See [Z, Section 3]).

1.2 Derived Picard Groups of Finite Dimensional Hereditary Algebras

This subsection is a summary of [MY] by J. Miyachi and A. Yekutieli.

First note that we work with right modules, hence the definition of Riedtmann quiver differs from that of [Hap] and [MY]. (They work with left modules.)

Let A be a finite dimensional k -algebra. A complex $T \in D^b(\text{Mod-}A \otimes_k A^{\text{op}})$ is called a *two-sided tilting complex* if there exists another complex $T^\vee \in D^b(\text{Mod-}A \otimes_k A^{\text{op}})$ such that $T \otimes_A^{\mathbf{L}} T^\vee \cong T^\vee \otimes_A^{\mathbf{L}} T \cong A$. The *derived Picard group* of A (relative to k) is

$$\text{DPic}_k(A) := \{\text{two-sided tilting complexes}\} / \text{isomorphisms}$$

with the identity element A , product $(T_1, T_2) \mapsto T_1 \otimes_A^{\mathbf{L}} T_2$ and inverse $T \mapsto T^\vee := \mathbb{R} \text{Hom}_A(T, A)$.

A tilting complex induces an equivalence of triangulated categories,

$$- \otimes_A^{\mathbf{L}} T : D^b(\text{Mod-}A) \longrightarrow D^b(\text{Mod-}A).$$

See [Y], [MY] for more details.

Let $\vec{\Delta} = (\vec{\Delta}_0, \vec{\Delta}_1)$ be a finite quiver. The Riedtmann quiver $\vec{\mathbb{Z}}\vec{\Delta}$ of $\vec{\Delta}$ is defined as follows : The set of vertices $(\vec{\mathbb{Z}}\vec{\Delta})_0$ is given by $\mathbb{Z} \times \vec{\Delta}_0$. Given an arrow $a : x \rightarrow y$ in $\vec{\Delta}$ and an integer n , there are arrows $(n, a) : (n, y) \rightarrow (n, x)$ and $(n, a)^* : (n, x) \rightarrow (n+1, y)$. The translation τ and the polarization μ on $\vec{\mathbb{Z}}\vec{\Delta}$ are defined by $\tau(n, x) = (n-1, x)$ and $\mu(n, a) = (n-1, a)^*$, $\mu((n, a)^*) = (n, a)$. For an arrow $\alpha : \xi \rightarrow \eta$ in $\vec{\mathbb{Z}}\vec{\Delta}$, we have $\mu(\alpha) : \tau(\eta) \rightarrow \xi$.

The *path category* $k\langle \vec{\mathbb{Z}}\vec{\Delta} \rangle$ (in the sense of [MY]) is the category whose set of objects is $(\vec{\mathbb{Z}}\vec{\Delta})_0$, morphisms are generated by identities and the arrows and the only relations arise from incomposability of paths.

Let η be a vertex of $\overrightarrow{\mathbb{Z}\Delta}$, let $\xi_1, \dots, \xi_p$ be complete representatives of the set $\{\xi \mid \text{there is an arrow } \alpha : \xi \rightarrow \eta\}$ and let $\{\alpha_{ij}\}_{j=1}^{d_i}$ be the set of arrows from ξ_i to η . The *mesh* ending at η is the subquiver of $\overrightarrow{\mathbb{Z}\Delta}$ with vertex $\{\eta, \mu(\eta), \xi_1, \dots, \xi_p\}$ and arrows $\{\alpha_{ij}, \mu(\alpha_{ij}) \mid i = 1, \dots, p, j = 1, \dots, d_i\}$. The *mesh ideal* in the path category $k\langle \overrightarrow{\mathbb{Z}\Delta} \rangle$ is the ideal generated by the elements

$$\sum_{i=1}^p \sum_{j=1}^{d_i} \alpha_{ij} \circ \mu(\alpha_{ij}) \in \text{Hom}_{k\langle \overrightarrow{\mathbb{Z}\Delta} \rangle}(\mu(\eta), \eta).$$

The *mesh category* $k\langle \overrightarrow{\mathbb{Z}\Delta}, I_m \rangle$ is defined as the quotient category of the path category modulo mesh ideal.

Let $A = k\overrightarrow{\Delta}$ be the path algebra of $\overrightarrow{\Delta}$ and let $\mathbf{mod}\text{-}A$ denote the category of finite right A -modules. The k -dual $A^* = \text{Hom}_k(A, k)$ of A is a two-sided tilting complex. We write $\tau_A \in \text{DPic}_k(A)$ for the element represented by $A^*[-1]$. Let us agree that τ_A also denotes the autoequivalence $-\otimes_A^{\mathbf{L}} A^*[-1]$ of $D^b(\mathbf{mod}\text{-}A)$.

Let P_x be the indecomposable projective right A -module corresponding to a vertex $x \in \overrightarrow{\Delta}_0$. Define $\mathcal{B} \subset D^b(\mathbf{mod}\text{-}A)$ to be the full subcategory with objects $\{\tau_A^n P_x \mid x \in \overrightarrow{\Delta}_0, n \in \mathbb{Z}\}$.

The following theorem which is important in [MY] is also important for us.

Theorem 1.5 ([MY, Theorem 2.6]). *There is a k -linear equivalence*

$$G : k\langle \overrightarrow{\mathbb{Z}\Delta}, I_m \rangle \longrightarrow \mathcal{B}, \quad G(n, x) = \tau_A^{-n} P_x.$$

The equivalence G sends the mesh ending at $\eta = (n, y)$ to the exact triangle called *Auslander-Reiten triangle*

$$(1) \quad \tau_A(G(\eta)) = G(\tau(\eta)) \xrightarrow{\oplus G(\beta_{ij})} \bigoplus_{i=1}^p \bigoplus_{j=1}^{d_i} G(\xi_i) \xrightarrow{\oplus G(\alpha_{ij})} G(\eta) \xrightarrow{[1]}$$

where $\beta_{ij} = \mu(\alpha_{ij})$. By the definition of the path category the sets of arrows $\{\alpha_{ij}\}_{j=1}^{d_i}$ and $\{\beta_{ij}\}_{j=1}^{d_i}$ are basis of $\text{Hom}_{k\langle \overrightarrow{\mathbb{Z}\Delta}, I_m \rangle}(\xi_i, \eta)$ and $\text{Hom}_{k\langle \overrightarrow{\mathbb{Z}\Delta}, I_m \rangle}(\tau(\eta), \xi_i)$ respectively. Fix isomorphisms

$$\begin{aligned} \text{Hom}_{k\langle \overrightarrow{\mathbb{Z}\Delta}, I_m \rangle}(\xi_i, \eta) &\cong \text{Hom}_{k\langle \overrightarrow{\mathbb{Z}\Delta}, I_m \rangle}(\tau(\eta), \xi_i)^* \\ \alpha_{ij} &\mapsto \beta_{ij}^* \end{aligned}$$

where $\{\beta_{ij}^*\}_{j=1}^{d_i}$ is the dual basis of $\{\beta_{ij}\}_{j=1}^{d_i}$. Set $V_i = \text{Hom}_{k\langle \overrightarrow{\mathbb{Z}\Delta}, I_m \rangle}(\xi_i, \eta) \cong \text{Hom}_{k\langle \overrightarrow{\mathbb{Z}\Delta}, I_m \rangle}(\tau(\eta), \xi_i)^*$. Then there are canonical morphisms

$$(2) \quad \varphi_i : V_i \otimes_k G(\xi_i) \longrightarrow G(\eta), \quad \psi_i : \tau_A(G(\eta)) \longrightarrow V_i \otimes_k G(\xi_i).$$

Then the Auslander-Reiten triangle (1) has the form

$$(3) \quad \tau_A(G(\eta)) \xrightarrow{\oplus \psi_i} \bigoplus_{i=1}^p V_i \otimes_k G(\xi_i) \xrightarrow{\oplus \varphi_i} G(\eta) \xrightarrow{[1]}.$$

Let $\text{Aut}((\overrightarrow{\mathbb{Z}} \overrightarrow{\Delta})_0)$ be the permutation group of the vertex set $(\overrightarrow{\mathbb{Z}} \overrightarrow{\Delta})_0$ and let $\text{Aut}((\overrightarrow{\mathbb{Z}} \overrightarrow{\Delta})_0; d)^{(\tau)}$ be the subgroup of permutations which preserve arrow-multiplicities and which commute with τ , namely

$$\begin{aligned} \text{Aut}((\overrightarrow{\mathbb{Z}} \overrightarrow{\Delta})_0; d)^{(\tau)} = \\ \{ \pi \in \text{Aut}(\overrightarrow{\mathbb{Z}} \overrightarrow{\Delta})_0 \mid d(x, y) = d(\pi(x), \pi(y)) \text{ for all } x, y \in (\overrightarrow{\mathbb{Z}} \overrightarrow{\Delta})_0, \text{ and } \pi\tau = \tau\pi \} \end{aligned}$$

where $d(x, y)$ denotes the arrow-multiplicity from x to y .

Now we can state one of the main result of [MY], which shall be used to prove our main theorem.

Theorem 1.6 ([MY, Theorem 3.8]). *Let $\overrightarrow{\Delta}$ be a finite quiver without oriented cycles and let $A = k\overrightarrow{\Delta}$ be the path algebra of $\overrightarrow{\Delta}$ over an algebraically closed field k . If A has infinite representation type then there is an isomorphism of groups*

$$\text{DPic}_k A \cong (\text{Aut}((\overrightarrow{\mathbb{Z}} \overrightarrow{\Delta})_0; d) \ltimes \text{Out}_k^0(A)) \times \mathbb{Z}$$

where $\text{Out}_k^0(A)$ denotes the identity component of the group of outer automorphisms $\text{Out}_k(A)$.

Remark 1.7. In [MY], the case of the finite representation type is also computed.

2 Proof of Theorem 0.1

From now on we assume that k is an algebraically closed field and $N \geq 2$.

Let A be the path algebra of the N -Kronecker quiver $\overrightarrow{\Omega}_N$ shown in introduction. We denote by P_0, P_1 the indecomposable projective right A -modules associated to vertices $0, 1$ of $\overrightarrow{\Omega}_N$, and S_0, S_1 the simple A modules associated to vertices $0, 1$ of $\overrightarrow{\Omega}_N$. Then $S_1 = P_1$ and P_0 is the projective cover of S_0 .

The Riedtmann quiver $\overrightarrow{\mathbb{Z}} \overrightarrow{\Omega}_N$ of the N -Kronecker quiver $\overrightarrow{\Omega}_N$ is shown in Figure 1 below. (In Figure 1 vertices $(-n, 1)$ and $(-n, 0)$ are replaced by $\tau_A^n P_1$ and $\tau_A^n P_0$ respectively.)




Figure 1: Riedtmann quiver $\overrightarrow{\mathbb{Z}} \overrightarrow{\Omega}_N$

It is easy to see that

$$\text{Aut}((\overrightarrow{\mathbb{Z}} \overrightarrow{\Omega}_N)_0; d) \cong \mathbb{Z}$$

and there is a generator ρ of $\text{Aut}((\overrightarrow{\mathbb{Z}} \overrightarrow{\Omega}_N)_0; d)$ such that $\rho(0, 1) = (0, 0)$, $\rho(0, 0) = (1, 1)$. This ρ satisfies the relation $\rho^{-2} = \tau$. By Theorem 1.6 there exists a two-sided tilting complex ρ_A such that $\rho_A^{-2} \cong \tau_A$ and $\rho_A P_1 \cong P_0$.

From now on we write $\rho = \rho_A$ and $\tau = \tau_A$.

For $M \in D^b(\text{mod-}A)$ we use the following notation

$$\rho^n M = M \otimes_A^{\mathbf{L}} \overbrace{\rho \otimes_A^{\mathbf{L}} \cdots \otimes_A^{\mathbf{L}} \rho}^n.$$

Since $P_0 \cong \rho P_1$ and $\rho^2 \cong \tau^{-1}$ it follows that $\tau^{-n} P_1 \cong \rho^{2n} P_1$, $\tau^{-n} P_0 \cong \rho^{2n+1} P_1$, the Riedtmann quiver $\overrightarrow{\mathbb{Z}} \overrightarrow{\Omega}_N$ has the form shown in Figure 2.




Figure 2: Riedtmann quiver $\overrightarrow{\mathbb{Z}} \overrightarrow{\Omega}_N$

Set $V = \text{Hom}(P_1, \rho P_1) \cong \text{Hom}(P_1, P_0)$. The set of arrows $x_1, \dots, x_N$ is a basis of V . The graded vector space $\bigoplus_{n \geq 0} \text{Hom}(P_1, \rho^n P_1)$ has a graded algebra structure as in the same way of Section 1.1.

By the mesh relation and Theorem 1.5 we have the following proposition.

Proposition 2.1. *There is an isomorphism of graded algebras*

$$\gamma : \bigoplus_{n \geq 0} \text{Hom}(P_1, \rho^n P_1) \longrightarrow k\langle X_1, \dots, X_N \rangle / \left(\sum_{i=1}^N X_i^2 \right)$$

which sends $x_i \in V$ to X_i for $i = 1, \dots, N$.

Fix isomorphisms $V \cong \text{Hom}_{D^b(\text{mod-}A)}(\rho^n P_1, \rho^{n+1} P_1)$ for $n \geq 0$ and identify V with its dual V^* by sending $x_i \mapsto x_i^*$. Then the canonical morphisms φ_i and ψ_i in (2) of Section 1.2 have the following forms:

$$\begin{aligned} \varphi : V \otimes_k \rho^n P_1 &\longrightarrow \rho^{n+1} P_1, \quad v \otimes p_n \mapsto v(p_n), \\ \psi : \rho^{n-1} P_1 &\longrightarrow V \otimes_k \rho^n P_1, \quad p_{n-1} \mapsto \sum_{i=1}^N x_i \otimes x_i(p_{n-1}). \end{aligned}$$

The Auslander-Reiten triangle (3) has the form

$$(\mathcal{S}_n) \quad \rho^{n-1} P_1 \xrightarrow{\psi} V \otimes_k \rho^n P_1 \xrightarrow{\varphi} \rho^{n+1} P_1 \xrightarrow{[1]}.$$

Set $R = k\langle X_1, \dots, X_N \rangle / (\sum_{i=1}^N X_i^2)$. We denote by R_n the degree n component of the graded ring $R = \bigoplus_{n \geq 0} R_n = k\langle X_1, \dots, X_N \rangle / (\sum_{i=1}^N X_i^2)$. The following lemma is easy to show.

Lemma 2.2. *For each $n \geq 1$, there is a following short exact sequence of k -vector spaces*

$$(4) \quad 0 \longrightarrow R_{n-1} \xrightarrow{\Psi} R_1 \otimes_k R_n \xrightarrow{\Phi} R_{n+1} \longrightarrow 0,$$

where the second and third morphisms are defined as follows:

$$\begin{aligned} \Phi : R_1 \otimes_k R_n \ni f \otimes_k r &\mapsto fr \in R_{n+1} \\ \Psi : R_{n-1} \ni r_{n-1} &\mapsto \sum_{i=1}^N X_i \otimes_k X_i r_{n-1} \in R_1 \otimes_k R_n. \end{aligned}$$

Let $\gamma_n : \text{Hom}(P_1, \rho^n P_1) \xrightarrow{\sim} R_n$ be the degree n part of the isomorphism γ of Proposition 2.1. Then we have the commutative diagram

$$\begin{array}{ccccc} \text{Hom}(P_1, \rho^{n-1} P_1) & \xrightarrow{\text{Hom}(P_1, \psi)} & V \otimes_k \text{Hom}(P_1, \rho^n P_1) \\ \gamma_{n-1} \downarrow \wr & & \gamma_1 \otimes \gamma_n \downarrow \wr \\ 0 \longrightarrow & R_{n-1} & \xrightarrow{\Psi} & R_1 \otimes_k R_n \end{array}$$

where the top row is obtained by applying $\text{Hom}(P_1, -)$ to the first two terms of the exact triangle $(\mathcal{S}_n)$ and the bottom row is the first three terms of the exact sequence (4) of Lemma 2.2. Note, since two vertical arrows in the diagram are isomorphisms, the injectivity of Ψ implies that $\text{Hom}(P_1, \psi)$ is injective.

Proposition 2.3. $\rho^n = \rho \otimes_A^{\mathbf{L}} \cdots \otimes_A^{\mathbf{L}} \rho$ is a pure module for $n \geq 0$, (i.e., $H^i(\rho^n) = 0$ for all $i \neq 0$).

Proof. Since $\rho^n \cong \rho^n P_1 \oplus \rho^{n+1} P_1$ and $\rho P_1 \cong P_0$ are pure modules, it suffices to show that $\rho^{n+1} P_1$ is pure for $n \geq 1$. By induction on n we may assume that $\rho^{n-1} P_1$ and $\rho^n P_1$ are pure. Then the long cohomology sequence of the Auslander-Reiten triangle $(\mathcal{S}_n)$ implies that $H^i(\rho^{n+1} P_1) = 0$ for $i \neq -1, 0$.

Note that $H^{-1}(\rho^{n+1}) = 0$ if and only if

$$\text{Ext}^{-1}(A, \rho^{n+1} P_1) \cong \text{Hom}(A, H^{-1}(\rho^{n+1} P_1)) = 0.$$

Since $A \cong P_1 \oplus \rho P_1$ and $\text{Ext}^{-1}(P_1, \rho^n P_1) \cong \text{Hom}(P_1, H^{-1}(\rho^n P_1)) = 0$ by the induction hypothesis, we have

$$\text{Ext}^{-1}(A, \rho^{n+1} P_1) \cong \text{Ext}^{-1}(P_1, \rho^{n+1} P_1) \oplus \text{Ext}^{-1}(P_1, \rho^n P_1) \cong \text{Ext}^{-1}(P_1, \rho^{n+1} P_1).$$

Thus we have only to show $\text{Ext}^{-1}(P_1, \rho^{n+1} P_1) \cong \text{Hom}(P_1, H^{-1}(\rho^{n+1} P_1)) = 0$.

There is an exact sequence of A -modules

$$0 \rightarrow H^{-1}(\rho^{n+1} P_1) \longrightarrow \rho^{n-1} P_1 \xrightarrow{\psi} V \otimes_k \rho^n P_1 \longrightarrow H^0(\rho^{n+1} P_1) \rightarrow 0.$$

Applying the exact functor $\text{Hom}(P_1, -)$ to the first four terms of this sequence yields the following exact sequence

$$0 \longrightarrow \text{Hom}(P_1, H^{-1}(\rho^{n+1}P_1)) \longrightarrow \text{Hom}(P_1, \rho^{n-1}P_1) \xrightarrow{\text{Hom}(P_1, \psi)} V \otimes_k \text{Hom}(P_1, \rho^n P_1).$$

The injectivity of $\text{Hom}(P_1, \psi)$ implies $\text{Hom}(P_1, H^{-1}(\rho^{n+1}P_1)) = 0$. $\square$

Corollary 2.4.

- (1) $\tau^{n+1}[1]$ is a pure module for every $n \geq 0$.
- (2) $\rho^{-n}S_0 \cong \rho^{-(n+1)}P_1[1]$ is a pure module for every $n \geq 0$.

Proof. (1) We have $\tau^{n+1}[1] \cong \tau^n \otimes_A^{\mathbf{L}} A^* \cong \mathbb{R} \text{Hom}(\tau^{-n}, A^*) \cong (\tau^{-n})^*$, using the property of the Serre functor $-\otimes_A^{\mathbf{L}} A^*$ for the third isomorphism. By Proposition 2.3 $\tau^{-n} \cong \rho^{2n}$ is pure for $n \geq 0$. Hence $\tau^{n+1}[1]$ is also pure for $n \geq 1$

(2) By the direct calculation we have

$$S_0 \cong P_0 \otimes_A^{\mathbf{L}} A^* \cong (\tau P_0)[1] \cong (\rho^{-1}P_1)[1].$$

Therefore $\rho^{-n}S_0 \cong \rho^{-(n+1)}P_1[1]$. $\rho^{-2n}S_0 \cong (\tau^{n+1}P_0)[1] \cong P_0 \otimes_A \tau^{n+1}[1]$ and $\rho^{-(2n+1)}S_0 \cong (\tau^{n+1}P_1)[1] \cong P_1 \otimes_A \tau^{n+1}[1]$ are pure by (1). $\square$

Remark 2.5. In this paper we assume $N \geq 2$. If $N = 1$ then it is known that $\tau^3 \cong A[-2]$ ([MY, Theorem 4.1]). Hence the above corollary fails.

The following lemma shall be used in the proof of the Lemma 2.10.

Lemma 2.6. The canonical morphism

$$c : \text{Hom}(\rho^{-(n+1)}S_0, \rho^{-n}S_0) \otimes_k \rho^{-(n+1)}S_0 \longrightarrow \rho^{-n}S_0$$

is surjective.

Proof. The canonical morphism c is the 1-shift of the last two terms of the Auslander-Reiten triangle

$$(\mathcal{S}_{-(n+2)}) \quad \rho^{-(n+3)}P_1 \xrightarrow{\psi} V \otimes_k \rho^{-(n+2)}P_1 \xrightarrow{\varphi} \rho^{-(n+1)}P_1 \xrightarrow{[1]}.$$

under the isomorphisms $\rho^{-n}S_0 \cong \rho^{-(n+1)}P_1[1]$ and $V \cong \text{Hom}(\rho^{-(n+1)}S_0, \rho^{-n}S_0)$. Therefore the cokernel of c is $H^0(\rho^{-(n+3)}P_1[2]) = 0$ by Corollary 2.4.(2). $\square$

Let us review the definition of a t-structure. A t-structure in a triangulated category $\mathcal{T}$ is a pair $(\mathcal{T}^{\leq 0}, \mathcal{T}^{\geq 0})$ of full subcategories which are closed under isomorphisms such that, for $\mathcal{T}^{\geq n} := (\mathcal{T}^{\geq 0})[-n]$, $n \in \mathbb{Z}$ and $\mathcal{T}^{\leq n} := (\mathcal{T}^{\leq 0})[-n]$, $n \in \mathbb{Z}$ the following three conditions are satisfied:

- (t1) For $X \in \mathcal{T}^{\leq 0}$ and $Y \in \mathcal{T}^{\geq 0}$ we have that $\text{Hom}_{\mathcal{T}}(X, Y) = 0$.

(t2) $\mathcal{T}^{\leq 0} \subset \mathcal{T}^{\leq 1}$ and $\mathcal{T}^{\geq 1} \subset \mathcal{T}^{\geq 0}$.

(t3) For $X \in \mathcal{T}$ there is an exact triangle $A \rightarrow X \rightarrow B \xrightarrow{[1]}$ such that $A \in \mathcal{T}^{\leq 0}$ and $B \in \mathcal{T}^{\geq 1}$.

The full subcategory $\mathcal{H} := \mathcal{T}^{\geq 0} \cap \mathcal{T}^{\leq 0}$ is called the *heart* of the t-structure $(\mathcal{T}^{\geq 0}, \mathcal{T}^{\leq 0})$. The important fact is that the heart $\mathcal{H}$ of a t-structure is an abelian category. (See [GM, IV.4])

Let X be a projective variety over k and $\mathcal{O}_X(1)$ be an ample line bundle on X . Let $D^{\geq 0}(\text{coh } X)$ (resp. $D^{\leq 0}(\text{coh } X)$) be the full subcategory of $D^b(\text{coh } X)$ with objects $\mathcal{F}^\bullet$ such that $H^i(\mathcal{F}^\bullet) = 0$ for $i < 0$ (resp. $i > 0$). Then the pair $(D^{\geq 0}(\text{coh } X), D^{\leq 0}(\text{coh } X))$ is a t-structure in $D^b(\text{coh } X)$ and its heart is $\text{coh } X$. By Serre's vanishing theorem [Har, Proposition III.5.3], $D^{\geq 0}(\text{coh } X)$ (resp. $D^{\leq 0}(\text{coh } X)$) is a full subcategory with objects $\mathcal{F}^\bullet$ which satisfy

$$\begin{aligned} \mathbb{R} \text{Hom}^\bullet(\mathcal{O}_X, \mathcal{F}^\bullet(n)) &\in D^{\geq 0}(k\text{-vect}) \quad \text{for } n \gg 0 \\ (\text{resp. } \mathbb{R} \text{Hom}^\bullet(\mathcal{O}_X, \mathcal{F}^\bullet(n)) &\in D^{\leq 0}(k\text{-vect}) \quad \text{for } n \gg 0) \end{aligned}$$

where we denote

$$\mathcal{F}^\bullet(n) = \mathcal{F}^\bullet \otimes_X^{\mathbf{L}} \mathcal{O}_X(n) = \mathcal{F}^\bullet \otimes_X^{\mathbf{L}} \overbrace{\mathcal{O}_X(1) \otimes_X^{\mathbf{L}} \cdots \otimes_X^{\mathbf{L}} \mathcal{O}_X(1)}^n.$$

As we expect that the pair (P_1, ρ) is ample, reversing Serre's vanishing theorem, we define the following.

Definition 2.7. *The full subcategory $D^{\rho, \geq 0}$ (resp. $D^{\rho, \leq 0}$) of $D^b(\text{mod-}A)$ consists of objects $M^\bullet$ which satisfy*

$$\begin{aligned} \mathbb{R} \text{Hom}^\bullet(P_1, \rho^n M^\bullet) &\in D^{\geq 0}(k\text{-vect}) \quad \text{for } n \gg 0 \\ (\text{resp. } \mathbb{R} \text{Hom}^\bullet(P_1, \rho^n M^\bullet) &\in D^{\leq 0}(k\text{-vect}) \quad \text{for } n \gg 0). \end{aligned}$$

Remark 2.8. *Since $A \cong P_1 \oplus \rho P_1$, $M^\bullet \in D^{\rho, \geq 0}$ (resp. $D^{\rho, \leq 0}$) if and only if $\rho^n M^\bullet \in D^{\geq 0}(\text{mod-}A)$ (resp. $D^{\leq 0}(\text{mod-}A)$) for $n \gg 0$.*

Proposition 2.9. *The pair of full subcategories $(D^{\rho, \geq 0}, D^{\rho, \leq 0})$ is a t-structure in $D^b(\text{mod-}A)$.*

Proof. Since the other conditions of t-structure are obvious by Remark 2.8, it suffices to check the condition (t3), namely, for every complex $M^\bullet \in D^b(\text{mod-}A)$ there is an exact triangle

$$A^\bullet \longrightarrow M^\bullet \longrightarrow B^\bullet \xrightarrow{[1]}$$

such that $A^\bullet \in D^{\rho, \leq 0}$ and $B^\bullet \in D^{\rho, \geq 1} = D^{\rho, \geq 0}[-1]$.

By [Hap, Lemma I.5.2] an indecomposable object of $D^b(\text{mod-}A)$ is of the form $M[-i]$, where $M \in \text{mod-}A$ and $i \in \mathbb{Z}$. By Lemma 2.3, $\rho^n M[-i] = M \otimes_A^{\mathbf{L}} \rho^n[-i] \in D^{[i-1, i]}(\text{mod-}A)$. Hence $M[-i] \in D^{\rho, \geq 1}$ for $i \geq 2$ and $M[-i] \in D^{\rho, \leq 0}$ for $i \leq 0$. The case when $i = -1$ is reduced to the following lemma. $\square$

Lemma 2.10. *For every $M \in \mathbf{mod}\text{-}A$ there exists a submodule M' such that $\mathrm{Hom}(P_1, \rho^n M') = 0$ for $n \gg 0$ and $\mathrm{Ext}_A^{-1}(P_1, \rho^n M'') = 0$ for $n \gg 0$, where we set $M'' = M/M'$.*

Proof. Define $\mathcal{T}_n = \{N \in \mathbf{mod}\text{-}A \mid N \text{ is generated by } \rho^{-n}S_0\}$ and $\mathcal{F}_n = \{N \in \mathbf{mod}\text{-}A \mid \mathrm{Hom}_A(\rho^{-n}S_0, N) = 0\}$. Since $\mathrm{Ext}_A^1(\rho^{-n}S_0, \rho^{-n}S_0) \cong \mathrm{Ext}_A^1(P_1, P_1) = 0$ and A is hereditary, by [Hap, Lemma III.4.2], the pair $(\mathcal{T}_n, \mathcal{F}_n)$ is a torsion theory on the abelian category $\mathbf{mod}\text{-}A$. So, if we set $t_n(M), f_n(M)$ to be the image and cokernel of the canonical morphism

$$\mathrm{Hom}(\rho^{-n}S_0, M) \otimes_k \rho^{-n}S_0 \rightarrow M,$$

then $t_n(M) \in \mathcal{T}_n, f_n(M) \in \mathcal{F}_n$. By Lemma 2.6 the canonical morphism

$$\mathrm{Hom}(\rho^{-(n+1)}S_0, \rho^{-n}S_0) \otimes_k \rho^{-(n+1)}S_0 \longrightarrow \rho^{-n}S_0$$

is surjective, hence $t_n(M) \subset t_{n+1}(M)$. Since $\dim_k M < \infty$, there is an integer n_0 such that $t_n(M) = t_{n_0}(M)$ for all $n \geq n_0$. Then by definition $f_n(M) = f_{n_0}(M)$ for all $n \geq n_0$. Set $M' = t_{n_0}(M), M'' = f_{n_0}(M)$. Then $M' \in \mathcal{T}_n, M'' \in \mathcal{F}_n$ for all $n \geq n_0$.

It is easy to see that $\mathrm{Hom}(P_1, \rho^n N) \cong \mathrm{Ext}_A^1(\rho^{-(n-1)}S_0, N) = 0$ for $N \in \mathcal{T}_{n-1}$ and $\mathrm{Ext}_A^{-1}(P_1, \rho^n N) \cong \mathrm{Hom}_A(\rho^{-(n-1)}S_0, N) = 0$ for $N \in \mathcal{F}_{n-1}$. Therefore $\mathrm{Hom}(P_0, \rho^n M') = 0, \mathrm{Ext}_A^{-1}(P_0, \rho^n M'') = 0$ for all $n > n_0$. $\square$

Remark 2.11. *Set $\mathcal{T}' = \{N \in \mathbf{mod}\text{-}A \mid \mathrm{Hom}(P_1, \rho^n N) = 0 \text{ for } n \gg 0\}$ and $\mathcal{F}' = \{N \in \mathbf{mod}\text{-}A \mid \mathrm{Ext}_A^{-1}(P_1, \rho^n N) = 0 \text{ for } n \gg 0\}$. Then by the Lemma 2.10 it is easy to see that $(\mathcal{T}', \mathcal{F}')$ is a torsion pair on $\mathbf{mod}\text{-}A$.*

From this torsion pair we can define a t -structure in $D^b(\mathbf{mod}\text{-}A)$ by setting

$$\begin{aligned} D'^{\geq 0} &:= \{M \in D^{\geq 0}(\mathbf{mod}\text{-}A) \mid H^0(M) \in \mathcal{F}'\} \\ D'^{\leq 0} &:= \{M \in D^{\leq 1}(\mathbf{mod}\text{-}A) \mid H^1(M) \in \mathcal{T}'\}. \end{aligned}$$

(See [HRS, Proposition I.2.1]). *However, this is not a new t -structure. It is easy to see that $(D'^{\geq 0}, D'^{\leq 0}) = (D^{\rho, \geq 0}, D^{\rho, \leq 0})$.*

Let $\mathcal{H}^\rho$ be the heart of the t -structure $(D^{\rho, \geq 0}, D^{\rho, \leq 0})$. By Proposition 2.3, $P_0, P_1 \in \mathcal{H}^\rho$. Now let us consider the triple $(\mathcal{H}^\rho, P_1, \rho)$.

Proposition 2.12. *The pair (P_1, ρ) is ample on $(\mathcal{H}^\rho, P_1, \rho)$.*

Proof. We check the conditions (1) and (2) of Definition 1.1.

First note that the cokernel of the morphism $f : M \rightarrow N$ in the abelian category $\mathcal{H}^\rho$ is $\tau_{\geq 0}^\rho(\mathrm{Cone}(f))$, where $\tau_{\geq 0}^\rho : D^b(\mathbf{mod}\text{-}A) \rightarrow D^{\rho, \geq 0}$ is the truncation functor (See [GM, IV.4]). So f is surjective in $\mathcal{H}^\rho$ if and only if $\mathrm{Cone}(f) \in D^{\rho, \leq -1}$.

(1) It is easy to see that for every $M \in \mathbf{mod}\text{-}A$ there is an exact sequence

$$0 \rightarrow \mathrm{Hom}_A(P_1, M) \otimes_k P_1 \rightarrow M \rightarrow \mathrm{Hom}_A(P_0, M) \otimes_k S_0 \rightarrow 0$$

which is functorial in M . So for every $M \in D^b(A\text{-mod})$ and for every n there is an exact triangle

$$\mathbb{R} \operatorname{Hom}(P_1, \rho^n M) \otimes_k \rho^{-n} P_1 \longrightarrow M \longrightarrow \mathbb{R} \operatorname{Hom}(P_0, \rho^{n-1} M) \otimes_k \rho^{-n} S_0 \xrightarrow{[1]}.$$

Since $\rho^{-n} P_1 \in D^{\rho, \leq n}$ for every integer n , $\rho^{-n} S_0 \cong \rho^{-(n+1)} P_1[1] \in D^{\rho, \leq -1}$.

Now assume that $M \in \mathcal{H}^\rho$ and take an integer n such that the complex $\mathbb{R} \operatorname{Hom}(P_1, \rho^{n-1} M)$ is pure. Then $\mathbb{R} \operatorname{Hom}(P_1, \rho^{n-1} M) \otimes_k \rho^{-n} S_0 \in D^{\rho, \leq -1}$.

(2) Let

$$M \xrightarrow{f} N \longrightarrow L \xrightarrow{[1]}$$

be a triangle such that $M, N \in \mathcal{H}^\rho$ and $L \in D^{\rho, \leq -1}$. Take an integer n such that $\operatorname{Hom}(P_1, \rho^n L) = 0$. Then the induced morphism $\operatorname{Hom}(P_1, \rho^n M) \longrightarrow \operatorname{Hom}(P_1, \rho^n N)$ is surjective. $\square$

Combining Proposition 2.1 and Proposition 2.12 and applying Theorem 1.2 we obtain the following proposition.

Proposition 2.13. *The triple $(\mathcal{H}^\rho, P_1, \rho)$ is equivalent to the noncommutative projective scheme $\operatorname{proj} R = (\operatorname{qgr} R, \overline{R}, (1))$. i.e., the functor*

$$\overline{\Gamma}_* : \mathcal{H}^\rho \longrightarrow \operatorname{qgr} R, \quad \overline{\Gamma}_*(M) = \bigoplus_{n \geq 0} \mathbb{R} \operatorname{Hom}(P_1, \rho^n M)$$

is an equivalence of categories, $\overline{\Gamma}_(P_1) = \overline{R}$ and $\overline{\Gamma}_* \circ \rho = (1) \circ \overline{\Gamma}_*$.*

Now we complete the proof of our main theorem.

The functor

$$\Gamma_* : \operatorname{mod-}A \longrightarrow \operatorname{Gr} R, \quad \Gamma_*(M) := \bigoplus_{n \geq 0} \operatorname{Hom}(P_1, M \otimes_A \rho^n)$$

is right exact. Let $\mathbf{L}\Gamma_*$ be the left derived functor of Γ_* . A finite projective A -module P is of the form $P \cong P_1^{\oplus n} \oplus P_0^{\oplus m}$ for some $n, m \in \mathbb{Z}_{\geq 0}$. Then $\Gamma_*(P) \cong R^{\oplus n} \oplus R(1)^{\oplus m}$ is a finite R -module. Since every complex $M \in D^b(\operatorname{mod-}A)$ is isomorphic to a bounded complex of finitely generated projective A -modules, $\mathbf{L}\Gamma_*(M)$ has cohomologies which are finitely presented. Thus there is an exact functor

$$\mathbf{L}\Gamma_* : D^b(\operatorname{mod-}A) \longrightarrow D_{\operatorname{gr} R}^b(\operatorname{Gr} R), \quad \mathbf{L}\Gamma_*(M) = \bigoplus_{n \geq 0} \operatorname{Hom}(P_1, \rho^n M).$$

Since the quotient functor $\pi : \operatorname{Gr} R \longrightarrow \operatorname{QGr} R$ is exact by [Pop, Theorem 4.3.8], this functor can be extended to the exact functor $\pi : D_{\operatorname{gr} R}^b(\operatorname{Gr} R) \longrightarrow D_{\operatorname{qgr} R}^b(\operatorname{QGr} R)$. Define $\mathbf{L}\overline{\Gamma}_* := \pi \circ \mathbf{L}\Gamma_*$.

The canonical functor $c : D^b(\operatorname{qgr} R) \longrightarrow D_{\operatorname{qgr} R}^b(\operatorname{QGr} R)$ is an equivalence by [BvB, Lemma 4.3.3]. Hence, to prove our main theorem, we have only to show that $\mathbf{L}\overline{\Gamma}_*$ is an equivalence.

There is a commutative diagram

$$(5) \quad \begin{array}{ccc} D^b(\text{mod-}A) & \xrightarrow{\mathbf{L}\overline{\Gamma}_*} & D_{\text{qgr } R}^b(\text{QGr } R) \\ i_A \uparrow & & \uparrow c \\ \mathcal{H}^\rho & \xrightarrow[\overline{\Gamma}_*]{\sim} \text{qgr } R \hookrightarrow D^b(\text{qgr } R) & \end{array}$$

where i_A, i_R are inclusions.

It is clear that $\text{qgr } R$ generates $D^b(\text{qgr } R)$, i.e., the minimal triangulated full subcategory of $D^b(\text{qgr } R)$ containing $\text{qgr } R$ is $D^b(\text{qgr } R)$ itself. Therefore the image $(\mathbf{L}\overline{\Gamma}_* \circ i_A)(\mathcal{H}^\rho)$ generates $D_{\text{qgr } R}^b(\text{QGr } R)$. Thus $\mathbf{L}\overline{\Gamma}_*$ is essentially surjective. To prove $\mathbf{L}\overline{\Gamma}_*$ is an equivalence it suffices to show that $\mathbf{L}\overline{\Gamma}_*$ is fully faithful.

Since every complex $M^\bullet \in D^b(\text{mod-}A)$ is obtained from P_0, P_1 by taking finite number of cones and shifts, the problem is reduced to the following lemma.

Lemma 2.14. *The map*

$$\text{Hom}_{D^b(\text{mod-}A)}(P_i, P_j[l]) \xrightarrow{\mathbf{L}\overline{\Gamma}_* P_i, P_j[l]} \text{Hom}_{D_{\text{qgr } R}^b(\text{QGr } R)}(\mathbf{L}\overline{\Gamma}_*(P_i), \mathbf{L}\overline{\Gamma}_*(P_j)[l])$$

is an isomorphism for every $i, j \in \{0, 1\}$ and $l \in \mathbb{Z}$.

Proof. For the case $l = 0$, by the commutativity of the diagram (5), the map $\mathbf{L}\overline{\Gamma}_* P_i, P_j$ is equal to

$$\begin{aligned} \text{Hom}_{D^b(\text{mod-}A)}(P_i, P_j) &\cong \text{Hom}_{\mathcal{H}^\rho}(P_i, P_j) \\ &\cong \text{Hom}_{\text{qgr } R}(\overline{\Gamma}_*(P_i), \overline{\Gamma}_*(P_j)) \\ &\cong \text{Hom}_{D_{\text{qgr } R}^b(\text{QGr } R)}(\mathbf{L}\overline{\Gamma}_*(P_i), \mathbf{L}\overline{\Gamma}_*(P_j)). \end{aligned}$$

Hence $\mathbf{L}\overline{\Gamma}_* P_i, P_j$ is an isomorphism.

By [Be2, Appendix] there exists an exact functor $F : D^b(\mathcal{H}^\rho) \rightarrow D^b(\text{mod-}A)$ and a commutative diagram

$$\begin{array}{ccc} D^b(\mathcal{H}^\rho) & \xrightarrow{F} & D^b(\text{mod-}A) \\ i_H \uparrow & \nearrow i_A & \\ \mathcal{H}^\rho & & \end{array}$$

where i_H, i_A are inclusions.

The restriction $\mathbf{L}\overline{\Gamma}_* \circ F|_{\mathcal{H}^\rho}$ to $\mathcal{H}^\rho$ is equal to $c \circ i_R \circ \overline{\Gamma}_*$ with the notation in the diagram (5). Hence $\mathbf{L}\overline{\Gamma}_* \circ F$ is an equivalence. Therefore, the map

$$\text{Hom}_{D^b(\text{mod-}A)}(M^\bullet, N^\bullet[l]) \xrightarrow{\mathbf{L}\overline{\Gamma}_* M^\bullet, N^\bullet[l]} \text{Hom}_{D_{\text{qgr } R}^b(\text{QGr } R)}(\mathbf{L}\overline{\Gamma}_*(M^\bullet), \mathbf{L}\overline{\Gamma}_*(N^\bullet)[l])$$

is surjective for $M^\bullet, N^\bullet \in D^b(\text{mod-}A)$ and $l \in \mathbb{Z}$. Now it is easy to prove the lemma because $\text{Hom}_{\text{mod-}A}(P_i, P_j[l]) = 0$ for every $i, j \in \{0, 1\}$ and $l \neq 0$. $\square$

Theorem 2.15 (Theorem 0.1). *Let k be an algebraically closed field, and $A = k\overrightarrow{\Omega}_N$, the path algebra of the N -Kronecker quiver $\overrightarrow{\Omega}_N$. Set moreover $R = k\langle X_1, \dots, X_N \rangle / (\sum_{i=1}^N X_i^2)$. Then, for $N \geq 2$, the functor*

$$\mathbf{L}\overline{\Gamma}_* : D^b(\text{mod-}A) \longrightarrow D_{\text{qgr } R}^b(\text{QGr } R)$$

is an equivalence of triangulated categories.

Remark 2.16. *From the argument above, $\{\overline{R}, \overline{R(1)}\}$ is a full strongly exceptional sequence, and by [Pop, Theorem 4.5.2] the abelian category $\text{QGr } R$ has enough injectives. Hence we can apply [Bo, Theorem 6.2] (and the following Remark) and prove that the functor*

$$\mathbb{R}\text{Hom}(T, -) : D^b(\text{qgr } R) \longrightarrow D^b(\text{mod-}A)$$

is an equivalence of triangulated categories where we set $T = \overline{R} \oplus \overline{R(1)}$.

Since $A \cong \text{End}(T)$, T has a natural left A -module structure. It is known that the functor $-\otimes_A^{\mathbf{L}} T$ is a quasi-inverse to the functor $\mathbb{R}\text{Hom}(T, -)$.

It is easy to see that there is a commutative diagram

$$\begin{array}{ccc} D^b(\text{mod-}A) & \xrightarrow{\mathbf{L}\overline{\Gamma}_*} & D_{\text{qgr } R}^b(\text{QGr } R) \\ & \searrow -\otimes_A^{\mathbf{L}} T & \uparrow c \wr \\ & & D^b(\text{qgr } R). \end{array}$$

Thus the functors $\mathbf{L}\overline{\Gamma}_$ and $-\otimes_A^{\mathbf{L}} T$ are essentially isomorphic.*

DEPARTMENT OF MATHEMATICS,
KYOTO UNIVERSITY, KYOTO 602-8502,
JAPAN
e-mail: minamoto@kusm.kyoto-u.ac.jp

References

- [AZ] M. Artin, and J.J. Zhang, Noncommutative projective schemes, *Advances in Math.* **109** (1994), pp. 228-287.
- [Be] A.A. Beilinson, Coherent sheaves on $\mathbb{P}^n$ and problems of linear algebras, *Func. Anal. Appl.* **12** (1978), pp. 214-216.
- [Be2] A.A. Beilinson, On the derived category of perverse sheaves, in *K-theory, arithmetic and geometry (Moscow 1984-1986)*, pp. 27-41, *Lecture Notes in Math.* **1289**, Springer, Berlin, 1987.
- [Bo] A.I. Bondal, Representations of associative algebras and coherent sheaves, *Izv. Akad. Nauk SSSR Ser. Mat.* **53** (1989), 25-44.

- [BvB] A. Bondal, and M. Van den Bergh, Generators and representability of functors in commutative and noncommutative geometry, *Mosc. Math. J.*, **3** (2003), no. 1, pp. 1-36.
- [GM] S.I. Gelfand, and Yu.I. Manin, *Methods of Homological Algebras*, (Second edition) Springer Monographs in Mathematica. Springer-Verlag, Berlin, 2003.
- [Hap] D. Happel, *Triangulated Categories in the Representation Theory of Finite-Dimensional Algebras*, London Math. Soc. Lecture Notes **119**, University Press, Cambridge, 1987.
- [HRS] D. Happel, I. Reiten, and S. Smalø, Tilting in abelian categories and quasitilted algebras, *Memoirs of the AMS*, vol. **575**, Amer. Math. Soc., 1996.
- [Har] R. Hartshorne, *Algebraic Geometry*, Graduated Texts in Mathematics, Springer-Verlag, 1977.
- [KR] M. Kontsevich, and A. Rosenberg, Noncommutative smooth spaces, *The Gelfand Seminars 1996-1999*, pp. 85-108, Gelfand Math Sem., Birkhäuser Boston, Boston, MA, 2000.
- [MY] J. Miyachi, and A. Yekutieli, Derived Picard groups of finite dimensional hereditary algebras, *Compositio Math.* **129** (2001), no. 3, pp. 341-368.
- [O] D. Orlov, Derived categories of coherent sheaves and triangulated categories of singularities, *math.AG/0503632*.
- [Po] A. Polishchuk, Noncommutative proj and coherent algebras, *Math. Res. Lett.*, **12** (2005), 1, pp. 63-74.
- [Pop] N. Popescu, *Abelian Categories with Applications to Rings and Modules*, L.M.S. Monographs, Academic Press, New York, 1973.
- [SvB] J.T. Stafford, and M. Van den Bergh, Noncommutative curves and noncommutative surfaces, *Bull. AMS* **38** (2001), pp. 171-216.
- [Y] A. Yekutieli, Dualizing complexes, Morita equivalence and the derived Picard group of a ring, *J. London Math Soc.* **60** (1999), pp. 723-746.
- [Z] J. J. Zhang, Twisted graded algebras and equivalences of graded categories, *Proc. London math. soc.* (3) **72** (1996), no. 2, pp. 281-311